



# Delaware 2001

Resident Characteristics*	STATE	LOCALE		
		Urban	Small Town	Isolated
Nursing Home Residents	4,000	3,017	913	**
Percent of Nursing Home Residents	100.00	75.43	22.83	
Percent Female	74.04	74.13	73.71	
Age: <25	0.30	0.13	0.88	
Age: 25-54	3.15	3.48	2.30	
Age: 55-64	4.70	5.07	3.18	
Age: 65-74	10.43	10.28	10.84	
Age: 75-84	32.23	32.62	30.89	
Age: 85-94	38.55	38.35	39.10	
Age: 95+	10.65	10.08	12.81	
American Indian/Alaskan Native	0.13	0.03	0.44	
Asian/Pacific Islander	0.38	0.46	0.11	
Black	17.15	18.50	12.38	
Hispanic	0.88	1.03	0.33	
White	81.47	79.97	86.75	
Lived Alone	31.81	31.03	34.77	
Lived with Others	51.71	46.74	60.06	
Lived in Other Facility	16.49	22.22	5.17	
Admitted from Private Home with No Services	21.13	23.21	13.51	
Admitted from Private Home with Services	8.15	8.42	9.20	
Admitted from Assisted Living	6.82	8.93	3.16	
Admitted from Other Nursing Home	13.23	13.27	15.52	
Admitted from Acute Care Hospital	45.01	38.52	56.32	
Admitted from Other Setting	5.66	7.65	2.30	
CPS: Intact or Borderline Intact	25.25	25.19	24.42	
CPS: Mild to Moderately Severely Impaired	48.88	49.98	46.11	
CPS: Severely or Very Severely Impaired	25.88	24.83	29.46	
ADL: Independent or Needs Supervision	13.58	13.79	12.27	
ADL: Limited Physical Assistance	12.25	12.86	9.97	
ADL: Extensive Assistance in Early Loss ADLs	19.78	19.39	21.14	
ADL: Extensive Assistance in Later Loss ADLs	9.70	9.88	9.42	
ADL: Totally Dependent in Some ADLs	25.03	24.56	26.29	
ADL: Totally Dependent in All Four ADLs	19.68	19.52	20.92	



## Delaware 2001 (continued)

	STATE	LOCALE		
		Urban	Small Town	Isolated
CHESS: No Instability	36.23	38.05	28.92	**
CHESS: Slightly Unstable	31.08	30.23	34.50	
CHESS: Mildly Unstable	18.10	17.90	19.06	
CHESS: Moderately Unstable	8.95	8.68	9.97	
CHESS: Highly Unstable	5.65	5.14	7.56	
Total Case-Mix Index (average)	0.89	0.89	0.91	
Nursing Case-Mix Index (average)	0.82	0.82	0.85	
<b>Quality Indicators</b>				
Depression without Therapy	7.73	8.00	6.87	
9+ Different Medications	45.43	43.42	50.82	
Bladder/Bowel Incontinence without Plan	32.52	29.06	39.91	
Indwelling Catheters	6.98	6.13	9.86	
Tube Feeding	9.33	9.51	9.09	
Antipsychotic Use without Conditions	18.77	18.49	19.42	
Antianxiety/Hypnotic Use	18.05	18.30	17.57	
Daily Physical Restraint	4.18	3.88	5.48	
Falls	18.80	18.00	21.14	
Behavioral Symptoms Affecting Others	25.08	24.74	26.83	
Symptoms of Depression	14.48	14.47	14.63	
Bladder/Bowel Incontinence	63.18	64.59	59.83	
Fecal Impaction	0.28	0.33	0.11	
Urinary Tract Infections	10.95	11.20	9.97	
Weight Loss	15.81	14.44	20.00	
Dehydration	0.95	0.60	2.19	
Bedfast Residents	4.98	4.24	7.45	
Little/No Activity	14.58	13.50	17.96	
Stage 1-4 Pressure Ulcer	18.77	19.22	16.98	

\* All table entries after the first row are percentages. The second row provides row percentages. Other percentages are column percentages. For Quality Indicators, entries are the percent of residents with the problem in facilities within the specific geographic setting.

\*\* Too few residents for data to be displayed.

