



New Jersey 2001

Resident Characteristics*	STATE	LOCALE		
		Urban	Large Town	Small Town
Nursing Home Residents	45,483	44,504	421	558
Percent of Nursing Home Residents	100.00	97.85	0.93	1.23
Percent Female	72.61	72.52	77.20	76.16
Age: <25	0.67	0.68	0.00	0.36
Age: 25-54	3.62	3.66	1.66	2.51
Age: 55-64	4.25	4.26	2.38	4.48
Age: 65-74	10.97	11.02	6.65	10.22
Age: 75-84	30.97	30.91	31.83	34.95
Age: 85-94	40.34	40.31	47.51	37.99
Age: 95+	9.18	9.17	9.98	9.50
American Indian/Alaskan Native	0.28	0.27	1.45	0.00
Asian/Pacific Islander	0.76	0.77	0.48	0.18
Black	12.38	12.44	9.93	9.32
Hispanic	3.59	3.65	0.97	0.72
White	82.99	82.86	87.17	89.78
Lived Alone	17.70	17.84	25.00	9.88
Lived with Others	74.02	73.71	65.38	90.12
Lived in Other Facility	8.28	8.45	9.62	0.00
Admitted from Private Home with No Services	17.79	17.83	11.54	16.67
Admitted from Private Home with Services	5.55	5.59	7.69	3.40
Admitted from Assisted Living	5.45	5.52	0.00	3.09
Admitted from Other Nursing Home	13.73	13.77	7.69	12.65
Admitted from Acute Care Hospital	50.92	50.65	65.38	61.11
Admitted from Other Setting	6.57	6.64	7.69	3.09
CPS: Intact or Borderline Intact	26.91	26.94	23.52	26.88
CPS: Mild to Moderately Severely Impaired	48.69	48.67	57.48	43.55
CPS: Severely or Very Severely Impaired	24.40	24.39	19.00	29.57
ADL: Independent or Needs Supervision	12.84	12.78	12.59	17.56
ADL: Limited Physical Assistance	13.96	14.00	13.54	11.29
ADL: Extensive Assistance in Early Loss ADLs	20.53	20.45	24.47	24.19
ADL: Extensive Assistance in Later Loss ADLs	9.03	9.04	11.40	6.81
ADL: Totally Dependent in Some ADLs	22.24	22.26	22.09	20.61
ADL: Totally Dependent in All Four ADLs	21.40	21.47	15.91	19.53



New Jersey 2001 (continued)

	STATE	LOCALE		
		Urban	Large Town	Small Town
CHES: No Instability	43.66	43.60	43.94	48.03
CHES: Slightly Unstable	25.53	25.57	24.94	22.76
CHES: Mildly Unstable	16.25	16.27	14.73	15.41
CHES: Moderately Unstable	9.65	9.64	11.40	9.50
CHES: Highly Unstable	4.91	4.92	4.99	4.30
Total Case-Mix Index (average)	0.92	0.92	0.86	0.87
Nursing Case-Mix Index (average)	0.83	0.84	0.79	0.78
Quality Indicators				
Depression without Therapy	5.39	5.38	8.33	3.95
9+ Different Medications	38.08	38.09	38.72	36.74
Bladder/Bowel Incontinence without Plan	43.68	43.38	58.16	53.57
Indwelling Catheters	9.72	9.74	7.84	9.50
Tube Feeding	10.79	10.85	4.75	10.75
Antipsychotic Use without Conditions	18.56	18.60	16.58	17.31
Antianxiety/Hypnotic Use	13.93	13.89	11.31	19.14
Daily Physical Restraint	5.29	5.34	3.80	2.51
Falls	15.26	15.24	13.78	17.74
Behavioral Symptoms Affecting Others	18.48	18.54	17.62	13.64
Symptoms of Depression	10.44	10.41	13.57	10.59
Bladder/Bowel Incontinence	59.61	59.58	61.26	60.45
Fecal Impaction	0.20	0.20	0.24	0.18
Urinary Tract Infections	8.39	8.40	7.36	8.06
Weight Loss	18.66	18.68	17.03	17.92
Dehydration	1.24	1.26	0.24	0.90
Bedfast Residents	5.16	5.18	4.99	3.76
Little/No Activity	14.26	14.24	14.05	15.62
Stage 1-4 Pressure Ulcer	28.87	28.89	31.96	25.41

* All table entries after the first row are percentages. The second row provides row percentages. Other percentages are column percentages. For Quality Indicators, entries are the percent of residents with the problem in facilities within the specific geographic setting.

